



CLINICAL CHARACTERISTICS COMPARISON OF THE TYPES OF PAROXYSMAL SUPRAVENTRICULAR TACHYCARDIA ATTACK BETWEEN YOUNG AND ELDERLY PATIENTS

^{1,2}Si Dung Chu, ¹Giang Song Tran, ³Minh Thi Tran, ¹Khanh Quoc Pham

¹Vietnam National Heart Institute, Bachmai Hospital, Vietnam

²Training and Education of Healthcare Activities Center, Bachmai Hospital, Vietnam

³Model Internal Department, Vietnam University of Traditional Medicine, Vietnam

Objectives: This study is carried out to describe types of clinical characteristic of paroxysmal supraventricular tachycardia attack (PSVT) between young and elderly patients who underwent cardiac electrophysiological test. **Methods:** This is a cross sectional descriptive study, and prospective study. 182 patients who were diagnosed with PSVT attack underwent cardiac electrophysiological test at Vietnam Heart Institute during 01/2014-05/2017. The patients were divided into two groups: Group I (n=93) is the young patients (< 60 years of age) and Group II (n=89) is a elderly patients (≥ 60 years of age). **Results:** Group I (young patients) accounted for 51.1% and Group II (elderly patients) made up 48.9%. The rate of PVST attack in female was 70.9% while that in male was 29.1% ($p > 0.05$); the mean age of elderly patients was higher than younger patients ($p < 0.0001$). The PVST attacks have moredate positive correlation with structure ($R = 0.355$) and hypertension ($R = 0.314$), there are differences between both groups in the structure and hypertension. The structure and hypertension rates of patients in elderly patients group were higher than those in young patients group ($p < 0.0001$), there is a strongly positive correlation between the structure and hypertension patients ($R = 0.966$, $p < 0.0001$). Characteristics of PVST attack are 68.3% of atrioventricular nodal reentrant (AVNRT) and 29.6% of atrioventricular reciprocating tachycardia (AVRT) attacks; The AVNRT and AVRT attack rates in female were 74.8% and 61.8%, respectively, whereas the AVNRT and AVRT attack rates in male were 25.2% and 28.2%, respectively ($p > 0.05$). **Conclusion:** PVST attacks were more common in female patients than in male patients, the likelihood of having AVNRT was higher compared to AVRT attacks, and the mean age of AVNRT and AVRT attack in the elderly patients was higher than that in young patients.

Keywords: Aging, clinical characteristics, PVST, young patients, elderly patients.

1.

Introduction

Paroxysmal supraventricular tachycardia (PSVT) attack is an abnormally of rapid heart rhythm, which has an electropathologic substrate on above the atrioventricular bundle, it's quite common in clinical practice, type of arrhythmias often appears suddenly and can be self-

extinguished. PSVT has frequency of about 0.23-3.0% [1], [2]. In the United States, according to data from the MESA study showed that the prevalence is 2.25 per 1.000 persons, the incidence of new case is 35 out of 100.000 person each year, and about 89,000 new cases each year [2]. In Vietnam, there are no comprehensive



statistics on morbidity. The PSVT attack can occur in patients of any ages, It's causing many uncomfortable symptoms such as palpitations, pulsation in the neck, fatigue, light headedness, sweating, tightness, chest pains, shortness of breath, dizziness and even syncope, ect in some cases, the PSVT attacks prolonged for many hours, or many days can causing hemodynamic disorders, or heart failure, increase the risk of falls (especially in the elderly) [2], [3]. PSVT attack is one of the rhythmias causing patients to have to go to the hospital, or emergency department. Therefore, the treatments is very necessary to prevent further complications of this arrhythmias, and the ablation by intervention is very good solution [2], [3].

The PSVT includes the most common is atrioventricular nodal reentrant (AVNRT), quite common is atrioventricular reciprocating tachycardia (AVRT) with Orthodontic AVRT attack is more common than Antidromic AVRT. Where AVNRT and AVRT are two types of rhythms depending on the AV node (the AV node is part or whole of the loop (of the ring on the back). This is meaningful treatment because then the drugs affecting the AV node will be able to cut off the attack [2], [3].

For decades, there have been many methods of treatments for PSVT attack that have been applied and also have been certainly effective. The treatment methods for cutting off paroxysmal supraventricular tachycardia attack such as the flute therapy for vagal nerve, drug therapy, drug, electric shock, pacemaker frequency, etc to the prevention of recurrent attacks with drugs. Many treatment rhythm drugs were born, developed and now scientists are continuing to research and develop new antiarrhythmic drugs that are more effective [4], [5]. However, these are only temporary treatments that do not completely resolve the origin of the rhythm. Nowadays, the method of electrocardiography and radiotherapy of arrhythmia with

radiofrequency wave energy is considered to be the most effective and most thorough treatment. Most of the PSVT attack can be completely eliminated [2], [5], [6]. Although, selecting specific treatments for this arrhythmias depends on the types of PSVT attacks, it is very important to identify rapid onset ventricular tachycardia because it will allow the doctor to select the appropriate treatment and treatment strategies. Prophylaxis is appropriate for each nature of the attack.

In the world there have been many studies on the characteristics of the PSVT attack [2], [5], [6]. However, very few authors mentioned in detail the characteristics of the PSVT attack of young and elderly people, ages relationship between aging and cardiovascular system [7-12]. Therefore, we conducted this study to explore the "characteristics of the paroxysmal supraventricular tachycardia attack between young and elderly patients" and "to findout some disorders of heart rhythm in young people and the elderly with characteristics of the paroxysmal supraventricular tachycardia attack through electrophysiological probe".

2. Materials and methods

Study design: Cross-sectional descriptive study, prospective study

Participantss: 182 patients who were diagnosed with characteristics of the paroxysmal supraventricular tachycardia attack should be diagnosed and treated with RF at the Vietnam Heart Institute, Bach Mai Hospital, Vietnam from 01/2014 to 05/2017. The patients were divided into two groups: Group I were young patients (patients under 60 years of age) & Group II were elderly patients (patients \geq 60 years of age) [13], [14], [15].

Describes the cross sectional descriptive study, proceeding in sequence analysis. Study at the Vietnam Heart Institute, Bach Mai Hospital. The research was conducted on patients who meet the selection criteria during January 2014 - May 2017.



Sample Size: Following formula can be used for calculation of sample size for comparison between two groups. The PSVT attack can occur

at any ages; Therefore, both AVNRT and AVRT are selected [16], [17].

$$\text{Sample size} = \frac{2 \times (Z_{\alpha/2} + Z_{\beta})^2 P (1-P)}{(p1 - p2)^2}$$

$Z_{\alpha/2} = Z_{0.05/2} = 1.96$ (From Z table) at error of 5% $Z_{\beta} = Z_{0.20} = 0.842$ (From Z table) at 80% power.

$p1-p2$: Difference in proportion of events in two groups = $(0.6 - 0.3) = 0.3$ [18] $P = \text{PSVT type prevalence} = [\text{Prevalence in the case of AVNRT group } (p1) + \text{Prevalence in the case of AVRT group } (p2)]/2 = (0.6+0.3)/2 = 0.45$

We selected $p1 = 0.6$ and $p2 = 0.3$ [18]

Calculator for the samples size = 43, the mean that researcher needs 43 patients per group. And then, the sample size for the two groups are 86 patients.

We selected 182 patients to be sure there are enough samples size for control, ensure for the sample size to comparison.

Statistical Analysis: Data is processed on a computer with IBM SPSS 21.0 software. Use T-test to compare 2 mean values, when squared compare 2 percentage. The p-value of 0.05 was considered statistically significant [17].

The correlation coefficient (R) show the strength and direction of a relationship between two variables ranging the between -1 and +1 and belong to values R as follow:

Values $R = 0$: No linear relationship;

Values $R = -1$ or $+1$: Perfect negative/positive linear relationship;

Values $R = (-0.3: 0)$ or $(0: 0.3)$: Weak negative/positive linear relationship; Values $R = (-0.7:$

$-0.3)$ or $[0.3: 0.7)$: Moderate negative/positive linear relationship;

Values $R = (-1:-0.7)$ or $[0.7:1)$: Strong negative/positive linear relationship [17], [19].

2.2

. Criteria for diagnosis of paroxysmal supraventricular tachycardia attack

Paroxysmal supraventricular tachycardia (PSVT) attack is episodes of fast heart rate, start attack in a part of the heart above the ventricle.

The electrocardiogram pattern which demonstrates the Heart rate of regular ranges are higher than 140 to 250 beats per minute, the narrow QRS complex (< 120 milliseconds), regular, hidden or inverted P waves (behind the QRS complex) [20], [22].

2.2.1. Criteria for the AVNRT

AVNRT: Retry caused by nodal pathways or tracts. There are two types as typical (Slow-fast) and atypical (Fast-Slow) [6-7], [21-25].

a/ Typical AVNRT

- Dual A-V nodal physiology
- Retrograde AV conduction is through the fast way, the retrograde atrial activation sequence during tachycardia in concentric



- VA interval during the tachycardia < 60 ms at His potential recording site and < 90 ms at high right atrium
- PV stimulation during tachycardia should not preexcite the atrium when delivered at a time when His bundle is refractory

b/ Atypical AVNRT

- A retrograde dual AV conduction in some patients
- Tachycardia induction depends on a critical A-V or His-atrial natural during retrograde slow pathway conduction
- Retrograde atrial activation sequence during tachycardia is concentric
- A-V natural during tachycardia < 60 ms at His potential site and > 90 ms at high right atrium
- PV stimulation during tachycardia should not reset the atrium when delivered at a time that His bundle is refractory.

2.2.2. Criteria for the AVRT

AVRT: Reentry caused by accessory pathways have two types which are orthodromic AVRT and antidromic AVRT attacks [6-7], [21-25].

a/ Orthodromic AVRT: RP interval < PR interval or RP interval > PR interval with a slowly conducting accessory pathway; retrograde P waves (leads DI, DII, DIII, aVF, V1); delta wave seen with normal sinus rhythm, not with tachycardia.

b/ Antidromic AVRT: Short RP interval < 100 msec; regular, wide QRS complex ≥ 120 msec; delta waves seen with normal sinus rhythm and tachycardia; concealed accessory pathways do not show delta waves.

3. Results

3.1. Clinical Characteristics:

182 patients who were diagnosed with paroxysmal supraventricular tachycardia attack underwent electrocardiography and radiofrequency

radiofrequency ablation tests, and were divided into two groups: Group I (younger group) and Group II (elderly group).

The mean of age is 53.1 ± 16.1 years [18: 78], they have 53 male of 182 patients (29.1%) and 129 female of 182 patients (70.9%) with $p = 0.344 (>0.05)$.

Results showed that group I (n=93) had 51.1% and mean age of 40.5 ± 12.6 years old [18:59], and group II (n = 89) had 48.9% and mean age of 66.3 ± 5.2 years old [60:78] ($p < 0.0001$). Group I had 30 males (32.3%) and 63 females (67.7%) with $p > 0.05$, group II had 25 males (25.8%) and 66 females (74.2%) with $p > 0.05$.

The most common symptom was palpitations found in 179/182 (98.4%), chest pain was found in 74/182 (40.7%), shortness of breath was found in 31/182 (17.0%), dizziness was found in 38/182 (20.9%), and 5/182 patients (2.7%) had fainting and syncope (all syncope patients were met in young group); Many patients have these symptoms combined.

There are 110 out of 182 patients (60.4%) with structurally normal of heart system. The structurally abnormal disease was found in 72 out of 186 patients (39.6%); Among 182 patients, there are 66 patients with hypertension (36.3%), 5 patients with diabetes, two patients with only diabetes (1.1%), three patients with Coronary artery disease (1.6%), two patients with mitral valve disease (1.1%), one patients with aortic valve disease (0.54%), and one patient with hypertrophic cardiomyopathy (0.54%). The distribution of the hypertension, only diabetes, coronary artery, mitral valve, aortic valve disease in each group is 20/66 (30.3%), 0%, 0%, 1/2 (50%), and 0%; and 46/66 (69.7%), 2/2 (100%), 3/3 (100%), 1/2 (50%), 1/1 (100%), respectively. Besides, there are 4 patients with Chronic obstructive pulmonary disease (COPD) of



which 2 patients were found in Group I and 2 patients in Group II.

The comparison between the two age groups and PSVT attacks have no difference with $p = 0.211$ (> 0.05). Comparison rate calculate between the two group and other types of PSVT attacks (AVNRT, AVRT attack) showed that no difference between the types of PSVT attacks and aging with $p > 0.05$ ($p = 0.211$, and $p = 0.211$, respectively).

The calculation rate for the structurally abnormal disease was 72/182 (39.6%) and the young and eldly group had 21/93 (22.6%) and 51/89 (57.3%), respectively; There is a significant difference between the structurally abnormal and two group ($p < 0.0001$), and there is moredate correlation between the

abnormally normal disease and aging with $R = 0.355$ (Correlation is significant at the 0.01).

The calculation rate for the hypertension disease was 66/182 (36.3%) in which goup I and group II had 20/93 (21.5%) and 46/89 (51.7%), respectively; There is a highly significant difference between the hypertension and two group ($p < 0.0001$), and there is moredate correlation between the hypertension disease and aging with $R = 0.314$ (Correlation is significant at the 0.01).

Besides, there is the strongly positive correlation between the structure and hypertension with $R = 0.966$, $p < 0.0001$ (Correlation is significant at the 0.01)

3.2. Characteristics of type of paroxysmal supraventricular tachycardia attack pacing

Table 1: Types of paroxysmal supraventricular tachycardia attack

Type attack \ Group	Group I (n=93)		Group II (n=89)		Aging (n=182)	
	n	%	n	%	n	%
AVNRT (n=127)	61	65.6%	66	74.2%	127	68.3%
R, P	R = - 0.093, P = 0.211				R = - 0.144, P = 0.052	
AVRT (n=55)	32	34.4%	23	25.8%	55	29,6%
R, P	R = 0.093, P = 0.211				R = - 0.093, P = 0.211	
Total	93	100%	89	100%	182	100%

Table 1: Group I and Group II were the most common in AVNRT attack with 61/93 (65.6%) and 66/89 (74.2%) with $p > 0.05$, respectively; while AVRT attack in these two groups was less frequent accounting for 34.4% (32/93 cases) and 25.8% (23/89 cases), respectively with $p > 0.05$. The distribution of the PSVT attack pacing between groups I and II was not significantly different ($p > 0.05$). There is a relationship between AVNRT/AVRT attack and aging, but with weak negative/positive correlation ($R = - 0.144$, and $R = 0.144$); Moreover, the different between AVNRT/AVRT attack and aging was not significant ($p > 0.05$).

Table 2: Gender distribution in all types of paroxysmal supraventricular tachycardia attack pacing

Type attack \ Group	Group I (n=93)		Group II (n=89)		Aging (n=182)	
	Male	Female	Male	Female	Male	Female
AVNRT (n=127)	18 (29.5%)	43 (70.5%)	14 (21.2%)	52 (78.8%)	32 (25.2%)	95 (74.8%)
R, P	R = 0.081, P = 0.439		R = 0.179, P = 0.093		R = 0.131, P = 0.077	
AVRT (n=55)	12(37.5%)	20(62.5%)	9 (39.1%)	14 (60.9%)	21 (38.2%)	34 (61.8%)



R, P	R = - 0.081, P = 0.439	R = - 0.179, P = 0.093	R = -0.131, P = 0.077
p	> 0.05	> 0.05	> 0.05

Table 2: Comparing in detail on the gender in each group showed that the female in each type attack (AVNRT and AVRT) was higher than the male in the each group I (70.5% and 29.5%; 62.5% and 37.5%, respectively) with very weak correlation, and Group II (78.8% and 21.2%; 61.8% and 38.2%) with weak correlation, it's no difference with $P > 0.05$.

Table 3: Mean age of all types of the PSVT attack pacing

Type of PSVT attack		Group	Group I ($x \pm SD$) <i>n</i> = 93	Group II ($x \pm SD$) <i>n</i> = 89	Total (<i>n</i> = 182)
1	AVNRT (<i>n</i> =127)		<i>n</i> = 61	<i>n</i> = 66	$P > 0.05$
	Mean of age		41.5 ± 11.6	66.8 ± 5.4	
	P		$P < 0.0001$		
2	AVRT (<i>n</i> =55)		<i>n</i> = 32	<i>n</i> = 23	
	Mean of age		38.6 ± 14.2	64.8 ± 4.1	
	P		$P < 0.0001$		
3	Total		40.5 ± 12.6	66.3 ± 5.2	
	P		$P < 0.0001$		

Table 3: In both groups (I and II), years of age in patients with AVNRT was higher than AVRT attack, the difference was not statistically significant with $p > 0.05$. Mean age of AVNRT and AVRT attack in two groups was different with $p < 0.0001$.

Overall, The female in AVNRT and AVRT were higher than male in both groups (The female with AVNRT and AVRT were 74.8% and 61.8%, respectively; The male with AVNRT and AVRT were 25.2% and 38.2%, respectively), the relationship between two gender's weak relationship, but it's also no difference with $p > 0.05$.

4. Discussion

4.1. Clinical Characteristics :

We were selected ages for two groups as WHO and many authors in the world did [14], [2627], Besides, some developed nations can selected two groups of age

by < 65 year and ≥ 65 year as Leonardo AO in the USA, studied about PSVT attacks in the general population by two groups < 65 years and ≥ 65 years [28].

The results showed that the PSVT attacks were quite common in middle-aged and elderly patients, and were also quite common with the mean age of 53.1 ± 16.1 years, which is lower than Chen SA's study [6] probably age of the Taiwanese life expectancy was higher than that of Vietnamese; although the mean age is the same, some authors such as Hiroko N (52 ± 14 years, *n* = 23) [21], the Japan has high life expectancy, Hiroki N only studied on type AVNRT attack and only report mean age of 23 patients with fast-slow AVNRT (range: 26 to 64 years) among 138 AVNRT patients Mean age in elderly group (66.5 ± 5.4 years) is the same as in study of



Peter MK (66.4 ± 1.7 years for the patients group ≥ 60 years) [11].

Although the rate of the female was higher than male in both groups, but the difference was not statistically significant ($p > 0.05$). Almost authors also showed that prevalence of gender in patients with PSVT attack had no difference between males and females. The rate of gender in Muhammad A (n=200) was 53.5% males and 46.5% females ($p < 0.05$) [20].

The palpitation symptom was the most common (98.4%), the same as other authors [17], Bottoni H showed that palpitation was absolutely found (100%) [7]; However, the study showed that 5 out of 182 patients with fainting and syncope (2.7%), Bottoni H showed that syncope was found in AVRT and AVNRT with respective rate of 8.7% and 8.3% [7]. This means that we must be aware of the possible dangers of this arrhythmias because the PSVT attacks can occur in patients who are associated with a variety of complex arrhythmias or complex myocardial structure [12]. Remarkably, two patients have syncope in the young group. There are no difference in the PSVT attacks and two groups ($p > 0.05$). According to many studies sudden onset can occur suddenly and at any age [15], [28].

There are difference between two groups and myocardial structure as well as hypertension disease with high significance ($p < 0.0001$). The relationship between two group and myocardial structure as well as hypertension disease are fairly strong relationship ($R = 0.355$ and $R = 0.314$, respectively), which mean that the aging has fairly strong relationship with structurally abnormal and hypertension disease [12], [15], [26], [29]. Bottoni found the hypertension disease of AVRT and AVNRT of 8% and 31% ($p < 0.01$), lower than our result because Bottoni had two groups with almost young patients including AVRT and AVNRT group with 25 ± 16 and 37 ± 17 years of age, respectively [7]. Yangni O

showed that the heart disease in patients of over 70 years was 35 out of 141 patients (25%) associating to the tachycardia attack [4]. Especially, the structure and hypertension were very strong relationship ($R = 0.966$, $p < 0.0001$).

4.2. Characteristics of the type of paroxysmal supraventricular tachycardia attack pacing

Type of paroxysmal supraventricular tachycardia attack in both young and elderly groups were predominant AVNRT attack, accounting for 65.6% and 74.2%, respectively, whereas AVRT attack found in these two groups was less frequent, just 32/93 (34.4%) and 23/89 (25.8%), respectively. There was no difference in the type of PSVT attack pacing between group I and II ($p > 0.05$).

Table 4: Comparison with some authors on distribution for type of SVT attacks

Authors	AVNRT (%)	AVRT (%)
Yangni O (n=141) [4]	73%	27%
Bottoni N (n=94) [7]	51.1%	48.9%
Joerg L (n= 395) [25]	78.0%	22.0%
Our research (n=182)	68.3%	29.6%



Table 4 are comparison amongs some authors in the terms of the distribution of types PSVT attack pacing. Our results showed that rate of AVNRT attack were higher than the author's Bottoni [7] and were lower than author's Yangni O and Joerg [4], the study of Yangni only focused on patients of older than 70 years [1] and Joerg L had higher mean of age [25]; While AVRT attack of our research was lower than author's Bottoni [7] and higher than author's Joerg L [25] and a littler higher than Yangni O [4]; This probaly because our research on both in young and elderly people, but the rate of older patients is quite high, and most of the authors above studied on all ages. Moreover, Yangni O only focused on rather older patients (> 70 years) [4]. The higher the age had increase in female/male rate. Elderly patients had higher possibility of AVNRT attacks than younger patients, but elderly patients had lower possibility of AVRT attack than younger patients. Patients with AVNRT attack had a higher proportion of female than male in both groups, but the difference was not statistically significant ($p > 0.05$).

In our study, the rate of AVNRT attack in group II was higher than the group I, this difference was not statistically significant ($p > 0.05$), whereas the rate of AVRT attack in group II was lower than group I, with $p > 0.05$ (Table 2). As the increase age of AVRT attacks is increasingly less, because the pathway is tends to decrease [5], [29], [30], however, the results are not clear between two groups ($p > 0.05$).

Relationship between age and gender in type of PSVT attack pacing: The male/female ratio between groups I and group II were different with $P > 0.05$; but the male/female ratio between AVNRT and AVRT groups was different with $P < 0.05$.

In both groups I and II, mean age of patient with AVNRT attack was significantly higher than AVRT but the difference was not statistically significant at $p > 0.05$.

Mean age between AVNRT and AVRT was significantly different with $p < 0.0001$.

5. Conclusion

The study of 182 patients with PSVT attack, Group I (young group) had 51.1% and Group II (elderly group) had 48.9%. Clinical characteristics and types of SVT attack as follow:

The PSVT attack was the most common found in female more than male ($p > 0.05$).

The mean age of elderly patients was higher than younger patients ($p < 0.0001$).

The PVST attacks have moderate correlation with structure and hypertension, both in the structure and hypertension in eldely patients were higher than young patients ($p < 0.0001$).

There is the strongly positive correlation between the structure and hypertension patients ($p < 0.0001$).

Charactetistics of PSVT attack with type AVNRT were more than AVRT attack. The elderly group with AVNRT, AVRT attacks with higher mean age than younger groups. The AVNRT and AVRT attack in female were than in male ($p > 0.05$).

The AVNRT attack was most common found in typical AVNRT from atypical AVNRT ($p < 0.0001$), the elderly patients with typical AVNRT attack was higher than typical AVNRT attack in young patients ($p > 0.05$).

Reference

1. Katritsis D.G., Camm A.J., Gersh B.J. (2016), *Clinical Cardiology: Current Practice Guideline*, Oxford University Press. 2016: 538.
2. Leonardo A.O., Humberto V., Frank D.S., et al. (1998), Paroxysmal Supraventricular Tachycardia in the General Population. *Journal of the American College of Cardiology*; 31 (1): 150-157.



3. Eugene B., Douglas P.Z., Peter L. (2003), *Heart Disease: A textbook of cardiovascular medicine*, 6th edition, W.B. Saunders Company, A Harcourt Health Sciences Company; 1 (1): 1-386.
4. Yangni N'Da' O., Brembilla-Perrot B (2008), Clinical characteristics and management of paroxysmal junctional tachycardia in the elderly. *Archives of Cardiovascular Diseases*, 101 (3): 143–148.
5. De Marneffe M., Jacobs P., Haardt R., et al. (1986), Variations of normal sinus node function in relation to age: role of autonomic influence, *Eur Heart J.*, 7 (8): 662-672.
6. Chen S.A., Chiang S.A., Yang C.J., et al. (1994), Accessory Pathway and Atrioventricular Node Reentrant Tachycardia in Elderly Patients: Clinical Features, Electrophysiologic Characteristics and Results of Radiofrequency Ablation, *J Am Coll Cardiol*; 23:702 – 708.
7. Bottoni N, Tomasi C, Donateo P, et al. (2003). Clinical and electrophysiological characteristics in patients with atrioventricular reentrant and atrioventricular nodal reentrant tachycardia, *Europace*, 5 (3): 225-229.
8. Karavidas A., Lazaros G., Tsiachris D. (2010), Aging and the Cardiovascular System, *Hellenic J Cardiol*; 51: 421-427.
9. North B.J., Sinclair D.A. (2012), The Intersection Between Aging and Cardiovascular Disease, *Circulation Research*; 110 (8): 1097-1108.
10. Chester J.G., Rudolph J.L. (2011), Vital Signs in Older Patients: Age-Related Changes, *J Am Med Dir Assoc.*; 12 (5): 337-343.
11. Peter M.K., Prashanthan S., Simon P.F., et al. (2004), Electrophysiologic and electroanatomic changes in the human atrium associated with age, *Journal of the American College of Cardiology*; 44 (1): 109-116.
12. Schwartz J.B. (1999), *Cardiovascular Function and Disease in the Elderly*, Northwestern University, 1999; 1-6.
13. Gürleyik G., Gürleyik E. (2003), Age-related clinical features in older patients with acute appendicitis, *Eur J Emerg Med.*; 10(3): 200-203.
14. WHO. (2002), *Health statistics and information systems*, Proposed working definition of an older person in Africa for the MDS Project.
15. Kistler P.M., Sanders P., Fynn S.P. (2004), Electrophysiologic and electroanatomic changes in the human atrium associated with age. *Journal of the American College of Cardiology*; 44 (1): 109116.
16. Jaykaran C., Tamoghna B. (2013), How to Calculate Sample Size for Different Study Designs in medical Research? *Indian J Psychol Med*; 35 (2): 121-126.
17. University of the West of England (2011), *Sample Size and Power in Clinical Trials*, North Bristol, NHS trust.
18. Randall AC., Mitchell J.S., Jay S. (2010), Common Types of Supraventricular Tachycardia: Diagnosis and Management. *Am Fam Physician*; 81 (8): 942-952.
19. Ratner B. (2009), The Correlation Coefficient: Its values range between +1/-1, or do they? *Journal of Targeting, Measurement and Analysis for Marketing*; 17 (2): 139-142.
20. Muhammad A., Abdul M., Amir H. (2007), Comparison of efficacy of intravenous adenosine and verapamil in acute paroxysmal



- supraventricular tachycardia in adults, *JSZMC*, 4 (30): 492-496.
21. Hiroko N., Naohito Y., Kenzo H. (1998), Heterogeneity of anterograde fast-pathway and retrograde slow-pathway conduction patterns in patients with the fast-slow form of atrioventricular nodal reentrant tachycardia: electrophysiologic and electrocardiographic considerations, *44th Annual Scientific Session of the American College of Cardiology, New Orleans, Louisiana*; S0735-1097 (98): 00433-1. 199.
 22. Braunwald Z.L. (2001), *Heart Disease: A textbook of cardiovascular medicine*, 6th Edition, W.B. Saunders Company: A Harcourt Health Sciences Company; 1 (25): 815-931.
 23. Demosthenes GK., Camm AJ. Classification and Differential Diagnosis of atrioventricular nodal re-entrant tachycardia. *EU Europace*; 8 (1): 29-36. (2006).
 24. Padanilam BJ., Manfredi JA., Steinberg LA., et al. (2008), Differentiating Junctional Tachycardia and Atrioventricular Node Re-Entry Tachycardia Based on Response to Atrial Extrastimulus Pacing, *Journal of the American College of Cardiology*; 52 (21): 1711-1717.
 25. Joerg L, Julia S., Ralph S. (2015), Electrophysiological studies in patients with paroxysmal supraventricular tachycardias but no electrocardiogram documentation: findings from a prospective registry, *EP Europace*; 17 (5): 801 – 806.
 26. Canonico V., De Caprio L, Vigorito C., et al. (1990), Differences in blood pressure profile between young and elderly hypertensive patients, *J Hum Hypertens*; 4 (4): 405-409.
 27. Ghnnam W.M. (2012), Elderly versus young patients with appendicitis 3 years experience. *Alexandria Journal of Medicine*; 48: 9-12. Rosen P., Barkin R.M. (2007), Supraventricular tachycardia, In. *5-Minute Emergency Medicine Consult*, 3th Edition, Wolters Kluwer Health: Lipincott Williams & Wilkins (2003): 1080 – 1081.
 28. Jennifer G.G., James L.R. (2011), Vital Signs in Older Patients: Age – Related Changes, *J Am Med Dir Assoc.*; 12 (5): 337-343.
 29. Shi X., Stevens G.H., Foresman B.H., et al. (1995), Autonomic nervous system control of the heart: endurance exercise training, *Med Sci Sports Exerc.*, 27: 1406-1413.
 30. Béatrice B.P., Arnaud O., Jean M.S. (2016), Influence of advancing age on clinical presentation, treatment efficacy and safety, and long-term outcome of pre-excitation syndromes: a retrospective cohort study of 961 patients included over a 25-year period, *BMJ Open*; 6(5): e010520.